

### **Acetaminophen (Tylenol)**

For use over 7 weeks of age. Other than for vaccine reaction, do not give acetaminophen to an infant under 3 months of age without consulting MD.

Give one dose every 4 hours as needed.

**Please be aware that there has been a change by the manufacturers. Products labeled as Infant Acetaminophen may be available as EITHER 80 mg/0.8 mL or 160 mg/5 mL. Please check the label of your medication to ensure that you are using the correct amount for the product that you have.**

Weight	Infant's concentrated 80mg/0.8mL	Infant's OR Children's oral suspension 160mg/5ml	Chewable tablets 80mg	Junior tablets 160mg
8 to 10 lbs	0.6 mL	2 mL		
11 to 13 lbs	0.8 mL	2.5 mL	1 tab	
14 to 16 lbs	1.0 mL	3.5 mL		
16 to 19 lbs	1.2 mL	4 mL	1 ½ tabs	
20 to 24 lbs	1.6 mL	1 tsp OR 5 mL	2 tabs	
25 to 29 lbs	2.0 mL	1 ¼ tsp OR 6.25 mL	2 ½ tabs	
30 to 34 lbs	2.4 mL	1 ½ tsp OR 7.5 mL	3 tabs	
35 to 40 lbs		1 ¾ tsp OR 8.75 mL		
41 to 49 lbs		2 tsp OR 10 mL	4 tabs	2 tabs
50 to 59 lbs		2 ½ tsp OR 12.5 mL	5 tabs	2 ½ tabs
60 to 69 lbs		3 tsp OR 15 mL	6 tabs	3 tabs
70 to 79 lbs		3 ½ tsp OR 17.5 mL		
Over 80 lbs		4 tsp OR 20 mL		4 tabs

### **Ibuprofen (Motrin or Advil)**

For use in infants and children over 6 months of age.

Give one dose every 6 hours as needed.

Weight	Infant drops 50mg/1.25mL	Children's oral suspension 100mg/5mL	Chewable tablets 50mg	Junior tablets 100mg	Adult tablets 200mg
12 to 16 lbs	1.25 mL	½ tsp OR 2.5 mL			
17 to 21 lbs	1.875 mL	¾ tsp OR 3.75 mL			
22 to 26 lbs	2.5 mL	1 tsp OR 5 mL	2 tabs		
27 to 29 lbs	3.125 mL	1 ¼ tsp OR 6.25 mL			
30 to 35 lbs	3.75 mL	1 ½ tsp OR 7.5 mL	3 tabs		
36 to 39 lbs	4.375 mL	1 ¾ tsp OR 8.75 mL			
40 to 49 lbs		2 tsp OR 10 mL	4 tabs	2 tabs	1 tab
50 to 54 lbs		2 ¼ tsp OR 11.25 mL			
55 to 58 lbs		2 ½ tsp OR 12.5 mL	5 tabs		
58 to 60 lbs		2 ¾ tsp OR 13.75 mL			
61 to 69 lbs		3 tsp OR 15 mL	6 tabs		
70 to 79 lbs		3 ½ tsp or 17.5 mL			
80 to 89 lbs		3 ¾ tsp or 18.7 mL			
Over 90 lbs		4 tsp OR 20 mL		4 tabs	2 tabs

***Diphenhydramine (Benadryl)***

For use in infants over 9 months of age, consult MD prior to use in any infant less than 9 months old. Give one dose every 6 hours as needed. For upper respiratory infections, one dose before bedtime is often enough for symptomatic relief.

<b>Weight</b>	<b>Children`s oral suspension 12.5mg/5mL</b>	<b>Chewable tablets 50mg</b>	<b>Swallow tablets 25 mg</b>
<b>10 to 14 lbs</b>	<b>½ tsp OR 2.5 mL</b>		
<b>15 to 19 lbs</b>	<b>¾ tsp OR 3.75 mL</b>		
<b>20 to 29 lbs</b>	<b>1 tsp OR 5mL</b>	<b>1 tab</b>	
<b>30 to 34 lbs</b>	<b>1 ¼ tsp OR 6.25 mL</b>		
<b>35 to 39 lbs</b>	<b>1 ½ tsp OR 6.25 mL</b>	<b>1 ½ tabs</b>	
<b>40 to 44 lbs</b>	<b>1 ¾ tsp OR 8.75 mL</b>		
<b>45 to 49 lbs</b>	<b>2 tsp OR 10 mL</b>	<b>2 tabs</b>	<b>1 tab</b>
<b>50 to 54 lbs</b>	<b>2 ¼ tsp OR 11.25 mL</b>		
<b>55 to 59 lbs</b>	<b>2 ½ tsp OR 12.5 mL</b>	<b>2 ½ tabs</b>	
<b>60 to 64 lbs</b>	<b>2 ¾ tsp OR 13.75 mL</b>		
<b>64 to 69 lbs</b>	<b>3 tsp OR 15 mL</b>	<b>3 tabs</b>	<b>1 ½ tabs</b>
<b>70 to 79 lbs</b>	<b>3 ¼ tsp OR 16.25 mL</b>		
<b>80 to 84 lbs</b>	<b>3 ½ tsp OR 17.5 mL</b>		
<b>84 to 89 lbs</b>	<b>3 ¾ tsp</b>		
<b>Over 90 lbs</b>	<b>4 tsp</b>	<b>4 tabs</b>	<b>2 tabs</b>